



Feral cat

Felis catus



A descendant of the African wild cat (*Felis silvestris lybica*), the common 'house' cat (*Felis catus*) has now been domesticated for about 4000 years. Although the domestic cat has a long history of association with humans, it retains a strong hunting instinct and can easily revert to a wild (feral) state when abandoned or having strayed from a domestic situation.

Feral cats live around dump sites, alleys or abandoned buildings, relying on humans by scavenging rubbish scraps and sheltering in abandoned structures. The true feral cat does not rely on humans at all, obtaining its food and shelter from the natural environment.

Legal requirements

The feral cat is a category 3, 4 and 6 restricted invasive animal under the *Biosecurity Act 2014*. This is a cat that is not owned. The feral cat must not be moved, fed, given away, sold, or released into the environment.

The Act requires everyone to take all reasonable and practical measures to minimise the biosecurity risks associated with invasive animals under their control. This is called a general biosecurity obligation (GBO).

At a local level, each local government must have a biosecurity plan that covers invasive animals in its area. This plan may include actions to be taken on feral cats. Some of these actions may be required under local laws. Contact your local government for more information.

Description

The feral cat differs little in appearance from its domestic counterpart; however, when in good condition, the feral cat displays increased overall muscle development, especially noticeable around the head, neck and shoulders, which gives the animal a more robust appearance. The average body weight of male feral cats is 3–6 kg, while females weigh 2–4 kg. Body weights vary with condition, with some extremely large specimens documented.

Australian feral cats are predominantly short-haired, with coat colours that range between ginger, tabby, tortoiseshell, grey and black. White markings may be present on the feet, belly, chest and throat; completely white feral cats are extremely rare. Coat colours are mostly the result of founding populations, although natural selection probably favours colours or patterns more suitable for each environment. The environment, predators and the ability to capture prey probably select coat colours to those that provide the most suitable camouflage and cause a predominance of these colours in subsequent offspring.

The feral cat is most active at night, with peak hunting activity occurring soon after sunset and in the early hours before sunrise. During the day it will rest in any number of den sites, which may include hollow logs, dense clumps of grass, piles of debris, rabbit burrows, and even the hollow limbs of standing trees.

The most obvious and characteristic field signs of feral cats are their scats (droppings). Unlike the domestic cat, the feral cat usually does not bury its scats, but leaves them exposed at prominent sites to warn other cats of its territorial boundary.

Life cycle

Male cats attain sexual maturity at about 12 months, whereas females are capable of reproduction at approximately seven months. Annually, and under ideal conditions, an adult female can produce up to three litters—each of usually four kittens, but varying from two to seven.

As the breeding instinct is triggered by the increasing length of daylight, litters are less frequent in winter. Most reproduction occurs during the spring and summer months, and is generally limited to two litters per year. Birth follows a gestation period of 65 days, and kittens may be reared in a single den site or may be frequently shifted to other sites within the female's home range. Family and litter bonding begin to break down when the kittens are approximately seven months old. The female's

ability to bear litters does not decrease with age, so reproduction continues for the course of her life.

Habitat and distribution

There is some evidence to suggest that the cat was present in Australia long before European settlement. This may have occurred as a result of Dutch shipwrecks and regular visits to northern Australia by early South-East Asian vessels as long as 500 years ago.

Post-settlement dispersal resulted from cats straying from areas of early colonisation. In the late 19th and early 20th centuries, large numbers of cats were purposely released in many rural areas to combat plague numbers of rabbits. Unwanted cats continue to be released into urban and rural areas by irresponsible pet owners.

The feral cat is now present Australia-wide, thriving under all climatic extremes and in vastly different types of terrain.

Feral cats maintain stable home ranges, the sizes of which depend upon the relative abundance of food and the availability of suitable den sites. Dominant male cats may have territories of up to 8 km², while the territories of females are smaller and may even be halved while kittens are being reared.

Scent glands are present on the chin, at the corners of the mouth, and in the anal region. Territorial boundaries are maintained by scent marking with the cheek glands, pole-clawing, urinating and leaving exposed faecal deposits. Although feral cats are often thought of as being solitary animals, studies show this behaviour is generally limited to hunting activities. At other times feral cats display a degree of social interaction that peaks during the breeding season. Group behaviour has been observed in wild populations, and it has been suggested that such behaviour is exhibited also in feral populations.

Groups usually comprise several related adult females, their young of both sexes, and an adult male—whose range may include other groups of females. Young females usually remain in a group, while young males either leave or are driven from the group as they reach sexual maturity.

Impacts

Effects on wildlife

The energy expended by an adult male cat requires it to consume 5–8% of its body weight in prey per day, while females raising kittens require 20%. Based on these figures, one study concluded that 375 feral cats on Macquarie Island would consume 56 000 rabbits and 58 000 sea birds per year. Where present on the mainland, rabbits may comprise up to 40% of a feral cat's diet. Cats are successful as a control mechanism only when rabbit densities are low. At other times cat predation does little to halt the build-up or spread of rabbit populations; rabbits merely help to support a larger number of cats.

When seasonal shortages of rabbits occur there is a corresponding rise in the number of native animals taken by cats.

The feral cat is an opportunistic predator, and dietary studies have shown that small mammals, birds, reptiles, amphibians, insects and even fish can be taken as prey.

Cat predation is particularly harmful in island situations, and a number of species have become extinct due to the introduction of cats. On the mainland, native animals—which already suffer due to the destruction of their habitats by man and other introduced animals—may be endangered further by cat predation. Actual competition for prey can cause a decline in the numbers of native predatory species such as quolls, eagles, hawks and reptiles. Not only do native animals bear the brunt of predation, but they also suffer the effects of a parasite that reproduces only in the intestine of the cat. This disease (toxoplasmosis) is particularly harmful to marsupials, which may develop blindness, respiratory disorders, paralysis, and suffer the loss of offspring through abortion and stillbirths.

Exotic disease—rabies

Due to their widespread distribution, feral cats may prove to be a major vector for this fatal viral disease if it ever enters Australia. Overseas studies have revealed that wounds inflicted by rabid cats are more dangerous than those caused by rabid dogs. While the bites of rabid dog are generally inflicted on the arms and legs, the cat attacks the head of its victim, biting and clawing viciously. These head and facial bites reduce the time taken for the virus to enter the central nervous system, lessening the chance of success from subsequent remedial treatment.

Control

Managing feral cats

The GBO requires a person to take reasonable and practical measures to minimise the biosecurity risks posed by feral cats. This fact sheet provides information and some options for controlling feral cats.

Exclusion fencing

Fencing is the only feasible method of control when special areas need total protection from cats. Feral cats have been successfully prevented from climbing over netted fences that use an electrified wire mounted 15 cm from the top and 10 cm outward from the fence. Non-electrified fencing should incorporate a netted ceiling, or a curved overhang, which prevents the cat from climbing straight up and over the fence.

Trapping

Rubber-jawed, leg-hold traps can be laid in the same manner as they are laid for dingoes and foxes. Leg-hold traps can help to capture true feral cats, which may avoid the live-capture box traps.

Ideal sites are those where territorial markers, such as faecal deposits and pole-clawing, are noticed. Tuna fish oil has shown some success as an attractant; however, feral cats seem more readily attracted to a site by some visual stimulus such as a bunch of bird feathers hung from a bush or stick.

Feral cats can be easily trapped in wire 'treadle-type' box traps (see diagram below). Attractants/lures include fresh meat or fish and should be placed so that they cannot be reached through the wire and be retrieved by clawing. A number of local governments hire cat traps for the purpose of removing stray and feral cats in urban situations.

Lures

Audible recorded lures for feral cats and other predators are available through a number of sources. These recordings mimic the distress call of a small animal and can be used to draw a predator to a bait or trap site. Visual lures (e.g. feathers) can also help attract cats.

Shooting


Night shooting is assisted by the cat's distinctive, green eyeshine under spotlight. Thermal imagers can also greatly assist in detecting cats at night. Cats have been successfully attracted by the use of a fox whistle.

Poisoning


There are limited viable options currently available to target cats with poison or poison baits in Queensland

More information

For more information contact your local government or visit biosecurity.qld.gov.au.



Rubber-jawed leg-hold trap



Treadle box trap



Fact sheets are available from biosecurity.qld.gov.au. The control methods recommended should be used in accordance with the restrictions (federal and state legislation, and local government laws) directly or indirectly related to each control method. These restrictions may prevent the use of one or more of the methods referred to, depending on individual circumstances. While every care is taken to ensure the accuracy of this information, the department does not invite reliance upon it, nor accept responsibility for any loss or damage caused by actions based on it.

